

The Welfare of Pigs

RSPCA Farm Animals Department *Information Sheet*

January 2009

Contents

- Introduction
- Pig natural history and behaviour
- Commercial pig production
- Key welfare issues
- Work of the RSPCA to improve pig welfare
- How you can help
- Recommended further information

How to contact us

Enquiries Service
RSPCA
Wilberforce Way
Southwater
Horsham
West Sussex
RH13 9RS

Tel:
+44 (0)300 1234 555

Web enquiry:
www.rspca.org.uk
(click on 'Contact Us')

Farm animals information:
www.rspca.org.uk/farmanimals

In an animal welfare emergency please contact the RSPCA's 24-hour cruelty and advice line:
+44 (0)300 1234 999

Introduction

Around half of the meat consumed in the world is pork, with over 150 million pigs reared annually in the EU for meat production. In the UK, around nine million pigs are slaughtered annually, whilst there are approximately 470,000 breeding pigs (sows, gilts and boars). A variety of housing and husbandry systems are in

use, including both indoor and outdoor production systems. Both can provide good living conditions, but some farming methods have ignored the pigs' needs by treating them as production units rather than fascinating and intelligent creatures with innate behavioural and physical needs.

Pig natural history and behaviour

Like all farm animals, modern farmed pigs have been bred from wild ancestors. Domesticated pigs are believed to be descendents of the native European wild boar (*Sus scrofa*). Until the 17th Century pigs in the UK were virtually pure descendents of the European wild boar. Such breeds were characterized by long, narrow bodies and long snouts, and strong rusty-coloured bristles. However, during the 18th and early 19th Centuries a number of pigs belonging to a markedly different type, which originated in China and South West Asia, were introduced into Britain. It is from a combination of these breeds that modern pigs/commercial breeds were developed. Wild boar still exist in the UK today, having been re-introduced on wild boar farms in the early 1990s. Several feral populations have since become established as a result of escapes/deliberate releases.

The main commercial pig breeds used in the UK are the Large White, Landrace and Welsh, although 75 per cent are cross-bred, being mixed with Duroc, Pietrain or Hampshire. Cross-breeds are thought to have better meat quality, be easier to handle, have a higher reproductive performance, and improved growth rate and carcass quality.

Before domestication, pigs were forest animals feeding on a variety of plants, insects and even small animals. A sow would produce one litter of three or four piglets a year and form a small social group with other sows. She would wean her piglets at around 12 weeks of age. Much of the day would be spent foraging or rooting for food and generally exploring the environment.

Although the modern pig is very different in appearance, it still shares most of its wild relative's instincts and behaviours.



Wild boar sow with boarlets.

Pigs are sensitive to extremes of climate and have no sweat glands (except for on the tip of the snout) nor thick hair cover, relying on fat for insulation. In cold weather pigs often huddle to keep warm, whilst in warm weather they wallow in water and mud to keep cool.

Pigs forage and root for food (a strong natural behaviour), and eat a wide range of vegetables and animal products, including carrion. In terms of senses, pigs rely on smell and hearing more than vision. They have a wide range of vocalisations and a good ability to locate odour sources.

Pigs are highly intelligent, inquisitive, adaptable, social animals that learn quickly; in many learning tests they can out-perform dogs.

Pigs prefer to live in stable families or small groups. However, they can be aggressive to each other, especially if unfamiliar animals are mixed. Boars are often solitary. Contrary to popular belief, pigs are

clean animals and if given properly designed living accommodation will always tend to use one particular area for dunging, thereby keeping their lying/sleeping area clean.

Commercial pig production

Approximately nine million pigs are slaughtered annually in the UK, whilst there are approximately 470,000 breeding pigs (sows, gilts and boars), although this number is falling. The average breeding herd consists of around 500 (indoors) to 800 (outdoors) individuals. The average rearing/finishing herd is around 1,200.

In recent years consumer demand for leaner meat has influenced the breeding, housing and management of pigs in order to produce quick growth and a low fat carcass. A result of this is that many of the famous old British breeds of pigs are no longer farmed commercially.

The age/weight at which pigs are slaughtered depends on the primary cut that is required. Pork pigs for small joints and fresh meat are slaughtered at about 55-62kg (15-16 weeks of age), whilst bacon pigs are slaughtered at around 90-100kg (approximately 22 weeks of age). However, there is currently a trend towards increasing slaughter weights. In contrast, an adult sow can weigh more than 300kg.

There are three main stages in pig production: accommodation for pregnant (dry) sows, for sows at the time of giving birth (farrowing) and suckling, and for growing the offspring for meat (finishing pigs). For each of these stages there are a variety of different housing systems in use. These vary from simple wooden huts to costly and elaborate controlled-environment buildings. The majority of pig units specialize in either breeding or fattening pigs, although some do both.

Breeding

Dry sow accommodation

Dry sows and gilts (i.e. young sows) are kept in a variety of housing systems, for example, individual stalls (outside the UK), groups with feeding stalls/electronic sow feeders, groups with food supplied on the floor or in a communal trough, and groups in fields, large yards or in family pen systems. Each housing system can vary with respect to the microclimate, type of flooring, use (or not) of straw or other bedding, diet, and frequency and method of feeding.



Outdoor breeding.

Around half of all the meat consumed in the world is pork, with over 150 million pigs being reared annually in the EU for meat production.

In recent years one of the most contentious issues has been the keeping of dry sows in individual, narrow stalls throughout their pregnancy. These sow stalls restrict movement as the sow cannot turn around, and has only limited movement forwards and backwards. A variation of this is the tether system in which the sow is attached to the floor by a short chain. Legislation banning the use of the sow stall and tether system was introduced in the UK from the end of 1998, whilst amendments to the EU Pig Husbandry Directive in 2001 meant that the use of tethers was banned in the rest of Europe from the beginning of 2006. The use of stalls throughout a sow's pregnancy, a system that is still in use throughout much of Europe and elsewhere, will be prohibited in the rest of Europe from 2013; however, their use will still be permitted for up to four weeks after service (mating).

The alternative to sow stalls, group housing, has the potential to provide a much better environment, giving more freedom to move, allowing social contact. It can lead to problems with aggression, particularly associated with feeding. One method used by some producers to help prevent aggression or bullying is a computer combined with an electronic sow feeder. Each sow wears a collar with a unique electronic ID, which allows individual recognition when the animal chooses to enter a box containing a feed dispenser. Exactly the right amount of food (as pre-programmed by the farmer) for that particular sow will then be delivered.

Up to 40 per cent of breeding sows in the UK are now kept outside under free-range conditions. Outdoor dry sows are kept in groups with access to large bedded huts that provide shelter and a dry lying area. Outdoor pigs have more space, and can explore their environment, forage and hence express their natural behaviours. However, if land or climatic conditions are unfavourable, welfare problems can result. Standards of management must be high to ensure adequate welfare standards are maintained.

Giving birth (farrowing)

Around 1-2 days before farrowing, the sow will attempt to build a nest, whether or not bedding material is available. In the wild, the sow would find a quiet area to farrow. Average farrowing duration is 2-3 hours depending on the age and experience of the sow and factors such as stress.

During farrowing, the sow lies on her side. At intervals between births she may change sides, stand up or sit down. Such movements can increase the risk of the sow crushing her piglets. Overlying/crushing can account for a number of piglet deaths prior to weaning. The greatest risk is when the sow is restless, and the incidence of overlying depends on the sow's previous experience, breed, strain, temperament, health, and the accommodation she is in.

The majority of sows in the UK (60 per cent +) farrow indoors in farrowing crates, into which they are placed up to

a week before they are expected to give birth. Farrowing crates are restrictive pens which prevent the sow from turning around, and which aim to reduce the risk of the sow lying on and crushing her newborn piglets. Farrowing crates have a 'creep area' with a heat lamp and food source to attract the piglets away from the sow when not suckling and so reduce over-lying/crushing.



A sow in a farrowing crate.

Up to 40 per cent of sows give birth to their litters outdoors in individual straw-bedded shelters called arcs from which they can come and go freely. The sows are moved into areas with access to these individual arcs prior to farrowing, where straw is provided which the sows will use to build a nest. Whilst the sow can come and go freely, movement of the piglets out of the arc is restricted for the first one to two weeks by a barrier ('fender') placed around the entrance. This ensures the piglets remain in the warmth of the arc during their most vulnerable phase, when they are more prone to chilling.

Key welfare issues

Inadequate stimulation (barren environments) and lack of space can worsen problem behaviours such as tail biting and aggression. Lameness is also a big problem, caused by joint disorders and/or infection, and foot injuries. The latter can be caused by factors including inappropriate flooring indoors and stony soils in outdoor systems.

Growing/finishing pigs

Tail docking

Approximately 80 per cent of piglets have their tails docked shortly after birth, the method used by many producers to try to reduce the likelihood of tail biting when pigs are older. Tail biting, a behavioural problem in pigs, can lead to serious injury and hence health and welfare problems. It is also a symptom of other welfare problems, indicating some form of physical or mental stress.

Whilst legislation prohibits the routine docking of piglets' tails, it can be carried out provided other measures have been taken first to try to reduce tail biting. Such measures

Regardless of whether the sows are kept indoors or outdoors, following weaning at around 21-28 days of age, the weaned pigs are moved into separate accommodation for growing/finishing, with groups of pigs of a similar age or size being kept together. The sow will be 'served' within 5 days of weaning, either via mating with a boar, or through artificial insemination (AI). On average, a sow will produce 2.5 litters per year.

Growing/finishing

Almost all pigs (96 per cent), including those born outdoors, are put into indoor accommodation for the growing/finishing period, although some systems provide an outdoor run together with an indoor area. Only 4 per cent of growing pigs will spend their entire lives outdoors. As with sow accommodation, a wide variety of systems are in use, from large straw barns housing 200 pigs, to kennel and run systems housing 20-25 pigs per pen.

Approximately 65 per cent of growing pigs are kept in indoor straw-based systems. However, many growing pigs are kept in barren, crowded pens with only slatted or concrete floors on which to live. Approximately 31 per cent of growing pigs are kept in slatted accommodation which, due to the system of manure (slurry) removal, do not allow the use of straw or similar manipulable material. Enriching the growing pigs' environment with some form of rooting material, such as straw, can help to reduce problems such as tail biting and may even lead to improved growth rates.

Most growing/finishing pigs are slaughtered between 18 and 30 weeks of age, depending on their commercial use (see above).

include the provision of environmental enrichment material (e.g. straw), reducing the stocking density, and increasing the amount of feeding space available.

Pigs have a strong instinct to root. However, legislation on provision of manipulable material is open to interpretation, resulting in variable requirements across the EU and often inadequate provisions for pigs.

Teeth clipping

Piglets may have their needle teeth clipped or ground to remove the sharp ends that can cause damage to the sows' teats and to other piglets. Both of these practices can cause pain as well as stress from being handled.

Castration

In the EU, it is legal to castrate male piglets without anaesthetic, provided it is done before seven days of age. However, this is no longer a common procedure in the UK, although it is still widely practiced elsewhere in Europe and beyond. It is estimated that 80 per cent of male piglets in the EU are castrated each year (approximately 100 million piglets).

Castration is carried out to reduce the risk of 'boar taint' in the resulting meat. Boar taint is an unpleasant flavour, sometimes noticeable to some consumers, in meat from

boars after they reach sexual maturity. In addition, castration reduces aggression in older pigs and makes managing them easier. However, in the UK, such issues are less of a problem as pigs are generally slaughtered at lighter weights, before they have reached sexual maturity.

Early weaning

Whilst the law requires a minimum weaning age of 28 days (21 days under certain provisos), the industry norm in the UK is 26 days. Early weaning of piglets can result in abnormal behaviours such as belly nosing and navel sucking.

Comparison of litters produced in 'natural' and commercial conditions.

	'Natural'	Commercial
Litter size (number of piglets)	6	12
Litter frequency (PA (number per year)	1	2.26
Weaning age	8 to 17 weeks	26 days*

*28 days in legislation, but can be 21 days with provisos

Lack of opportunity to express key natural behaviours

Pigs have a strong instinct to root. However, legislation on provision of manipulable material is open to interpretation, resulting in variable requirements across the EU and often inadequate provisions for pigs. Not only does this frustrate rooting behaviour but it can also cause injury and discomfort. Whilst 65 per cent of pigs in the UK are finished in straw-based systems the amount of straw provided varies. Some pigs will have access to a deep layer of straw covering most of the floor area, whilst others will be given an amount that is insufficient to allow them to carry out their desire to root.

Aggression

Aggression can occur in some systems, and can be sparked by factors such as mixing unfamiliar pigs, barren environments and inadequate feeding facilities. In addition, nursing sows with piglets can also be aggressive, as can boars.

Dry Sows

As previously discussed, sow stalls, whilst banned in the UK, are still permitted in the rest of the EU. These particular systems cause severe welfare problems as the sow is unable to turn around and most have no access to bedding; they are unable to take more than two steps forward or backward throughout most of their four-month pregnancy. In

Farrowing crates, whilst offering some protection from crushing for the first few days after farrowing, cause a large amount of stress to the sow, as a result of confinement.

contrast, when properly managed, designed and provided with bedding material, indoor group housing systems for breeding sows can provide very good welfare

conditions. Bedding material provides comfort and an opportunity for pigs to root and forage. This also helps to prevent boredom and stress which, in barren concrete or slatted floor enclosures with high stocking densities, can result in outbreaks of fighting, as well as increasing the risk of problem behaviours such as tail, vulva and ear biting.



Outdoor sow exhibiting nest building behaviour.

Farrowing sows and piglets

There are a number of welfare issues related to farrowing, affecting both the sow and her piglets.

Piglet mortality

In the UK around 13 per cent of piglets fail to survive to weaning. Mortality can be due to a number of factors such as chilling, starvation, disease, over-lying/crushing and cannibalism by the sow.

Farrowing crates

Farrowing crates, whilst offering some protection from crushing for the first few days after farrowing, cause a large amount of stress to the sow, as a result of confinement. The inability of the sow to build a nest, due to confinement and a lack of bedding materials, further frustrates her, as she is not able to perform this natural behaviour.



The RSPCA has funded research into ways to avoid the need for nose ringing in outdoor systems.

Nose ringing

Outdoor sows are often nose ringed in an attempt to prevent excessive rooting of the ground. If such rooting occurs on the soil floor of farrowing arcs, it can lead to increased mortality of the piglets through crushing, or through chilling as farrowing arcs are placed on uneven land leading to

draughts. As well as pain when the ring is inserted and the stress of handling, this practice leads to chronic pain; nose rings work by the action of discomfort and pain as the sow goes to root with her snout, causing her to stop. Evidence has shown that adding bulky feeds to improve satiety can help to reduce rooting, whilst the use of a 'sacrificial' area of the paddock with root crops added can direct sows' rooting to that particular area, reducing rooting within the farrowing area.

Management issues

Other problems can also result on outdoor units, as a result of inadequate levels of stock-keeper input and skill, inappropriate site suitability (soil type, weather conditions), inadequate shade/shelter provision, restricted access to

food and water (particularly in winter), accidents to/predation of piglets, and potentially higher levels of some diseases (though others are less common than on indoor units).

Transport

Pigs are notoriously poor travellers, and can suffer from travel sickness. They also have problems regulating their body temperature, and it is therefore recommended that pigs be fasted for a limited period prior to travel. A few pigs have a so-called 'stress' gene (the Halothane gene) which makes them even more susceptible to the effects of stress, such as at the time of loading/unloading and transport, although this has now been bred out of most commercial strains of pig.

Work of the RSPCA to improve pig welfare

Welfare standards

The RSPCA encourages pig producers to adopt the 'RSPCA welfare standards for pigs', which have been developed to ensure that higher standards of animal welfare are met at all stages of the animals' lives.

In the RSPCA welfare standards, conventional farrowing crate systems are not permitted; all farrowing accommodation must allow sows to turn around easily from five days after farrowing. We believe that confining sows for more than the first few days cannot be justified, given the little benefit of the crate, in terms of reduced crushing of piglets, after the first few days following farrowing. We would therefore like to see producers investigating alternatives, and phasing out conventional farrowing systems that involve close confinement of the sow for several weeks.

Teeth clipping, nose ringing and tail docking are not permitted except in exceptional circumstances. Even then, the herd veterinary surgeon and the farmer must write in to the RSPCA, setting out what other measures have been implemented to avoid carrying out the practice. In addition, certain other standards must be met before permission will be granted.

The RSPCA welfare standards require all pigs to be provided with environmental enrichment, in the form of straw or other suitable material.



RSPCA welfare standards for pigs.

Campaigning work

In autumn 2000, we joined with Eurogroup for Animals (an umbrella organisation representing welfare groups across the EU) to launch a campaign to improve EU law on pig husbandry. The RSPCA pressed for improvements to the EU Directive, including a ban on sow stalls and conventional farrowing crate systems, provision of a solid lying area and bedding/rooting material for all pigs, and better training for pig stock-keepers. In June 2001, the EU Agriculture Ministers agreed in principle to ban sow stalls – though not until 2013 – and to make provision of rooting material to all pigs compulsory, though the wording in the resulting law has led to significant variation in interpretation of the requirement, and many pigs are still not provided with suitable materials. Additional improvements for growing pigs such as increases in space allowance must await further discussions. The Directive is due to be reviewed in 2009 and it is hoped that some of these outstanding issues will be addressed.

Advocacy work

RSPCA Farm Animals Department staff represent the RSPCA on a number of government and industry committees, including the BPEX Pig Health and Welfare Council. Staff have also been invited to participate in a series of workshops on the issue of piglet castration, which are part of a large EU project. The project is looking at the attitudes of people from different countries to surgical castration without anaesthetic, current practices and alternatives, such as immunocastration or leaving males entire (uncastrated), and their implications.

Work to improve pig meat labelling

The RSPCA has, in conjunction with the pig industry, been developing a set of definitions to describe the various methods of pig production, with a view to using these as labels on pig meat products. Unlike the situation in relation to methods of egg and chicken production, there is currently no legal definition of terms such as 'free range' with regard to pig production. It is not surprising that there is confusion amongst consumers as to the production method of the pig meat they are purchasing. We hope that these definitions,

which cover terms such as 'free range', 'outdoor reared', 'outdoor bred' etc., will be used by retailers and become incorporated into their labelling of pork products. We believe that it is vital that consumers are provided with clear, transparent and consistent information in order to enable them to make informed purchasing decisions. Longer-term, the RSPCA would like to see the development and implementation of legislation on labelling of pig meat.

Research projects

The RSPCA is involved in a number of research projects, both in terms of providing funding, and/or as members of project steering groups.

Reducing tail biting in pigs

We are co-funding a collaborative three-year project, involving leading animal welfare scientists from the universities of Bristol and Newcastle, along with input from the UK pig industry itself. The project was initiated in 2006 to look at the problem of tail biting. The aim is to make use of the existing scientific and practical knowledge on tail biting and its causes by developing an on-farm husbandry advisory tool to assess the risk factors specific to individual farms and to produce farm-specific programmes to reduce the risk of tail biting outbreaks occurring.

In 2007 the RSPCA sponsored a workshop bringing together leading pig scientists, industry representatives and vets to discuss the issues and consolidate existing knowledge. We are also funding the development of the intervention advisory tool to be used in the event of an outbreak occurring.



The RSPCA takes every opportunity to try improve the lives of all pigs.

Recommended further information

- For further information, contact the RSPCA Farm Animals Department (contact details on page 1).
- Faucitano L and Schaefer AL (2008) Welfare of pigs from birth to slaughter. Wageningen Academic Publishers: Versailler, France.

Development of non-confinement, mutilation-free systems for pigs

The RSPCA is also funding a project being undertaken by the Oxford-based Food Animal Initiative, with the aim of developing a mutilation and confinement free system for pigs of all ages, that is commercially viable and could be applied by all pig producers, including those who currently keep their animals in standard indoor conditions.

In fact, the RSPCA has been seeking an answer to the farrowing crate dilemma for some time and over a number of years has funded research to develop and commercially test alternatives, which will protect piglets but still allow the sow to move freely. Several less restrictive designs are now being used on commercial farms in various countries including the UK, and results have been encouraging.

In addition to this work, a new Defra-funded project has begun at the University of Newcastle and the Scottish Agricultural College to look at developing and testing alternative farrowing systems. This three year project aims to use existing knowledge of the needs of both the sow and piglet to design a new farrowing system to optimise the health and welfare of both. A representative of the Farm Animals Department has been invited onto the project steering group.

Past projects we have initiated/funded have also included:

- Reducing aggression between pigs.
- Avoidance of nose-ringing in outdoor pigs.

How you can help!

If you eat pork, bacon or ham and are concerned about welfare, look out for products carrying the RSPCA's Freedom Food label. Freedom Food is the RSPCA's farm animal welfare and food labelling scheme that aims to ensure that animals are reared, handled, transported and slaughtered according to welfare standards developed and monitored by the RSPCA.

If more consumers insist on higher welfare products, more supermarkets will stock them, more farmers will be encouraged to improve their farming practices, and more farm animals will benefit. You may therefore wish to write to your local supermarket's head office and ask what standards of welfare the farms from which they source pig meat must meet.

You may also wish to express, in your own words, your concerns about the need to improve pig welfare to your local MP, and to:

The Rt Hon Hilary Benn MP
Secretary of State for Environment, Food & Rural Affairs
Department of Environment, Food & Rural Affairs
Nobel House
17 Smith Square
LONDON SW1P 3JR.